



US006855691B1

**(12) United States Patent
Stamler****(10) Patent No.: US 6,855,691 B1
(45) Date of Patent: Feb. 15, 2005****(54) METHODS FOR PRODUCING AND USING S-NITROSOHEMOGLOBINS****(75) Inventor: Jonathan S. Stamler**, Chapel Hill, NC (US)**(73) Assignee: Duke University**, Durham, NC (US)**(*) Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.**(21) Appl. No.: 08/616,371****(22) Filed: Mar. 15, 1996****Related U.S. Application Data****(60)** Provisional application No. 60/003,801, filed on Sep. 15, 1995.**(51) Int. Cl.⁷ A61K 38/42; C07K 14/805****(52) U.S. Cl. 514/6; 514/2; 514/832; 530/385; 530/829; 604/28****(58) Field of Search 514/2, 6, 832; 530/385, 829; 604/28, 52****(56) References Cited****U.S. PATENT DOCUMENTS**

4,900,719 A	2/1990	Means et al.	514/18
5,152,979 A *	10/1992	Hunter	424/78.38
5,380,758 A	1/1995	Stamler et al.	514/562
5,395,314 A *	3/1995	Klatz et al.	604/24
5,405,919 A	4/1995	Keefer et al.	525/377
5,427,797 A	6/1995	Frostell et al.	424/434
5,439,882 A *	8/1995	Feola et al.	514/6
5,480,866 A	1/1996	Bonaventura et al.	514/6
5,574,068 A	11/1996	Stamler et al.	514/562
5,593,876 A	1/1997	Stamler et al.	435/188
5,863,890 A	1/1999	Stamler et al.	
6,087,479 A	7/2000	Stamler et al.	530/363
6,255,277 B1	7/2001	Stamler et al.	
6,291,424 B1	9/2001	Stamler et al.	
6,583,113 B2 *	6/2003	Stamler et al.	514/6
2002/0052314 A1	5/2002	Stamler et al.	
2003/0007967 A1	1/2003	Stamler et al.	
2003/0022267 A1	1/2003	Stamler et al.	

FOREIGN PATENT DOCUMENTS

WO	WO 93/09806	5/1993
WO	WO 93/12068	6/1993
WO	WO 94/22306	10/1994
WO	WO 94/22482	10/1994
WO	WO 94/22499	10/1994
WO	WO 95/07691	3/1995
WO	WO 96/03139 A1	2/1996
WO	WO 96/15797	5/1996
WO	WO 96/16645	6/1996
WO	WO 96/17604	6/1996
WO	WO 96/30006	10/1996
WO	WO 97/18000	5/1997

OTHER PUBLICATIONS

60/003,801 (currently unavailable to the Examiner).*

U.S. Appl. No. 08/123,331, filed Sep. 17, 1993, now abandoned.

U.S. Appl. No. 08/438,418, filed May 10, 1995, now U.S. 6,255,277.

U.S. Appl. No. 08/460,465, filed Jun. 2, 1995, now U.S. 6,087,479.

U.S. Appl. No. 09/433,550, filed Nov. 4, 1999, now U.S. 6,174,539.

U.S. Appl. No. 09/621,610, filed Jul. 21, 2000, now U.S. 6,471,978.

U.S. Appl. No. 09/661,190, filed Sep. 13, 2000, now U.S. 6,352,709.

U.S. Appl. No. 07/791/668, filed Nov. 14, 1991, now abandoned.

U.S. Appl. No. 07/943,835, filed Sep. 14, 1992, now abandoned.

U.S. Appl. No. 08/198,854, filed Feb. 17, 1994, now abandoned.

U.S. Appl. No. 08/287,830, filed Aug. 9, 1994, now U.S. 5,593,876.

U.S. Appl. No. 08/437,868, filed May 9, 1995, now abandoned.

U.S. Appl. No. 08/907,217, filed Aug. 6, 1997, now U.S. 5,863,890.

U.S. Appl. No. 09/092,622, filed Jun. 5, 1998, now U.S. 6,291,424.

U.S. Appl. No. 09/835,038, filed Apr. 16, 2001, published as 20020052314 (cited on PTO form 1449). The status and currently pending claims on this application are not known to Applicant.

U.S. Appl. No. 10/216,865, filed Aug. 13, 2002, published as 20030007967 (cited on PTO form 1449). The status and currently pending claims of this application are not known to Applicant.

(List continued on next page.)

Primary Examiner—Bennett Celsa*(74) Attorney, Agent, or Firm*—Hamilton, Brook, Smith & Reynolds, P.C.**(57) ABSTRACT**

Nitric oxide (NO) interacts with hemoglobin (Hb) at its metal centers, whereas S-nitrosothiols (RSNOs) can donate the NO group to β 93 cysteine residues, thereby shielding the NO functionality from heme inactivation. S-nitrosylation of Hb is under the allosteric control of oxygen and the oxidation state of heme. NO group release from S-nitrosohemoglobin (SNO-Hb) is further facilitated by intracellular low molecular weight thiols, forming RSNOs which can be exported from the erythrocyte to regulate blood pressure. Hence, a dynamic cycle is established in which S-nitrosylation of Hb is initiated in the lung following oxygenation of red blood cells and is completed by SNO-Hb metabolism during arterial-venous transit. SNO-Hb can be formed by reaction of Hb with S-nitrosothiol. This procedure avoids oxidation of the heme. SNO-Hb in its various forms and combinations thereof (oxy, deoxy, met; S-nitrosylated to various extents) can be administered to a mammal in a method of therapy where it is desired to oxygenate, to scavenge free radicals, or to release NO groups to tissues. Thiols can also be administered to enhance the transfer of NO groups. Examples of conditions to be treated by SNO-Hb therapy include ischaemic injury, hypertension, angina, reperfusion injury and inflammations.

5 Claims, 6 Drawing Sheets